

Idempotents of the Hecke algebra become Schur functions in the skein of the annulus

Sascha G. Lukac

lukac@liv.ac.uk

Department of Mathematical Sciences, University of Liverpool,
Peach St, Liverpool L69 7ZL, England

Abstract

The Hecke algebra H_n contains well known idempotents E_λ which are indexed by Young diagrams with n cells. They were originally described by Gyoja [5].

A skein theoretical description of E_λ was given by Aiston and Morton [2]. The closure of E_λ becomes an element Q_λ of the skein of the annulus. In this skein, they are known to obey the same multiplication rule as the symmetric Schur functions s_λ as stated in theorem 8.2. But previous proofs of this fact as in [1] used results about quantum groups which were far beyond the scope of skein theory.

Our elementary proof was motivated by [6] and uses only skein theory and basic algebra.

1 The skein of a planar surface

We consider a planar surface F with designated n incoming and n outgoing boundary points for some integer $n \geq 0$. The Homfly skein $\mathcal{S}(F)$ is defined as the module of linear combinations of oriented tangles in F quotiented by regular isotopy (i.e. Reidemeister moves II and III), the two local skein relations in figure 1 where v and s are variables, and the relation that a disjoint simple closed curve can be removed from a diagram at the expense of multiplication with the scalar $\delta = \frac{v^{-1}-v}{s-s^{-1}}$. The last relation is a consequence of the other relations unless the remaining diagram is the empty diagram.

$$\begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} - \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = (s - s^{-1}) \cdot \left(\begin{array}{c} \downarrow \\ \downarrow \end{array} \right), \quad \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = v^{-1} \cdot \left(\begin{array}{c} \downarrow \\ \downarrow \end{array} \right)$$

Figure 1: Skein relations.

The scalars can be chosen as $\mathbb{Z}[v^{\pm 1}, s^{\pm 1}]$ with powers of $s - s^{-1}$ in the denominators. If necessary, we can choose the scalars to be the field of rational functions in v and s . This will occur in section 3 when we divide a quasi-idempotent by a scalar to turn it into an idempotent.

For the skein of any planar surface F there is a map $\rho : S(F) \rightarrow S(F)$ which is induced by switching all crossings and replacing s by s^{-1} and v by v^{-1} . The skein relations are preserved by this map. We remark that ρ is not linear.

2 Young diagrams and symmetric functions

2.1 Young diagrams

We consider partitions of non-negative integers where the summands are in non-strictly decreasing order. We consider partitions to be equal if they differ only by a number of zero-summands at the end.

A *Young diagram* is a graphical depiction of a partition. A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ becomes a flush-left arrangement of square cells with λ_1 cells in the first row, λ_2 cells in the second row, ... and λ_k cells in the k -th row.

We define $\lambda \subset \mu$ for partitions $\lambda = (\lambda_1, \dots, \lambda_k)$ and $\mu = (\mu_1, \dots, \mu_m)$ if $\lambda_i \leq \mu_i$ for all $1 \leq i \leq k$ (where $\mu_i = 0$ for $m < i \leq k$). This relation coincides in the graphical depiction with λ being a subset of μ .

We define $|\lambda|$ to be the number of cells of λ , i.e. for $\lambda = (\lambda_1, \dots, \lambda_k)$ we have $|\lambda| = \lambda_1 + \dots + \lambda_k$.

2.2 Symmetric functions

Following Macdonald [8], we consider the polynomial ring in variables $x_1, \dots, x_n$. For a partition $\lambda = (\lambda_1, \dots, \lambda_k)$ with $k \leq n$ we define

$$s_\lambda(x_1, \dots, x_n) = \frac{\det(x_i^{\lambda_j + n - j})_{1 \leq i, j \leq n}}{\det(x_i^{n - j})_{1 \leq i, j \leq n}}.$$

This quotient is a symmetric polynomial in $x_1, \dots, x_n$. In the inverse limit of the polynomial rings $\mathbb{Z}[x_1, \dots, x_n]$ for $n \rightarrow \infty$ we can define an element s_λ , called the λ -*Schur function* for any Young diagram λ . This element specializes to $s_\lambda(x_1, \dots, x_n)$ when we set $x_i = 0$ for all $i \geq n + 1$ provided that λ has at most n non-zero summands.

The most basic Young diagrams are on the one hand a row diagram with, say, n cells and on the other hand a column diagram with, say, k cells. One can check that $s_{(n)}$ is the n -th complete symmetric function h_n and that $s_{(1^k)}$ is the k -th elementary symmetric function e_k .

3 The skein R_n^n of a disk with $2n$ boundary points

We consider a rectangle F with n outgoing points at the top and n incoming points at the bottom, $n \geq 0$. We denote the Homfly skein of this surface by R_n^n . It is known to be isomorphic to the Hecke algebra H_n .

R_n^n gets an algebra structure by defining $D \cdot E$ as putting D above E so that the incoming points of D match with the outgoing points of E . Clearly, R_n^n is generated as an algebra by the elementary braids $\sigma_1, \dots, \sigma_{n-1}$ where σ_i agrees with the identity braid up to a single positive crossing between the arcs which connect the boundary points i and $i+1$ (numbered from the left).

As a module, R_n^n is linearly spanned by the positive permutation braids ω_π where π is a permutation on n letters (see [9] for details).

R_n^n has a well known quasi-idempotent a_n which is given in terms of the positive permutation braids ω_π on n strings as

$$a_n = \sum_{\pi \in S_n} s^{l(\pi)} \omega_\pi$$

where $l(\pi)$ is the writhe of the braid ω_π . This element ‘swallows’ an elementary braid σ_i at the expense of a scalar s (see [9]).

Lemma 3.1 $a_n \sigma_i = \sigma_i a_n = s a_n$ for any $1 \leq i \leq n-1$.

This behaviour implies that a_n is central in R_n^n because the elementary braids $\sigma_1, \dots, \sigma_{n-1}$ generate R_n^n . Lemma 3.1 enables an inductive proof that $a_n a_n = \alpha_n a_n$ where the scalar α_n is given by

$$\alpha_n = s^{\frac{n(n-1)}{2}} [n][n-1] \cdots [1]$$

where $[i] = \frac{s^i - s^{-i}}{s - s^{-1}}$ for any integer $i \geq 0$.

We recall the map ρ from section 1.

Lemma 3.2 $\rho\left(\frac{1}{\alpha_n} a_n\right) = \frac{1}{\alpha_n} a_n$ for any $n \geq 1$.

Proof We have $\rho(ab) = \rho(a)\rho(b)$ for any elements $a, b \in R_n^n$. This multiplicativity together with lemma 3.1 implies that

$$\sigma_i \rho(a_n) = \rho(\sigma_i^{-1}) \rho(a_n) = \rho(\sigma_i^{-1} a_n) = \rho(s^{-1} a_n) = s \rho(a_n)$$

in R_n^n . Hence, $a_n \rho(a_n) = \alpha_n \rho(a_n)$. When we apply the map ρ to this equation we get $\rho(a_n) a_n = \rho(\alpha_n) a_n$. The left hand sides of the previous two equations are equal because a_n is central. Hence, $\alpha_n \rho(a_n) = \rho(\alpha_n) a_n$ and thus $\rho\left(\frac{1}{\alpha_n} a_n\right) = \frac{1}{\alpha_n} a_n$. $\diamond$




Figure 2: The multiplication in the skein of the annulus $\mathcal{C}$.




Figure 3: The additional arcs for the closure map $\Delta_n : R_n^n \rightarrow \mathcal{C}$.

Aiston and Morton [2] constructed idempotents E_λ in R_n^n for any Young diagram λ with n cells. To do this, they considered a three-dimensional version of R_n^n where the distinguished boundary points are lined up along the cells of the Young diagram λ . In this setting they arranged several copies of idempotents a_i and variants b_i of them along the rows and columns of λ to get a quasi-idempotent. This element which is denoted e_λ in the equivalent R_n^n -setting satisfies $e_\lambda e_\lambda = \alpha_\lambda e_\lambda$. It is sufficient to prove that $\alpha_\lambda \neq 0$ for the specialization $v = s = 1$ which turns R_n^n into the symmetric-group algebra as explained in [1].

We consider the rational functions in v and s as the set of scalars for the skein R_n^n . Then, the element $E_\lambda = \frac{1}{\alpha_\lambda} e_\lambda$ is an idempotent of R_n^n .

4 The skein $\mathcal{C}$ of the annulus

The skein of the annulus shall be denoted by $\mathcal{C}$. Let D_1 and D_2 be two diagrams in the annulus $S^1 \times [0, 1]$. We can bring D_1 into $S^1 \times [0, 1/2)$, and D_2 into $S^1 \times (1/2, 1]$ by regular isotopies. Then the product of D_1 and D_2 is defined as the diagram $D_1 \cup D_2$. An example is shown in figure 2. The product is commutative since $D_1 D_2$ and $D_2 D_1$ differ by regular isotopy. The empty diagram is the identity.




Figure 4: Encircling a diagram in the annulus.

Figure 3 depicts an annulus with a set of n oriented arcs. A rectangle is removed from the annulus so that we can insert a diagram from R_n^n with matching orientations of the arcs. This factors to a map $\Delta_n : R_n^n \rightarrow \mathcal{C}$. This is a special case of a *wiring*. We abbreviate Δ_n by Δ if the index n is obvious from the context.

We denote by $\mathcal{C}_n$ the image of R_n^n under the closure map which is a submodule of $\mathcal{C}$. By $\mathcal{C}_+$ we denote the submodule of $\mathcal{C}$ spanned by all $\mathcal{C}_0, \mathcal{C}_1, \dots$

We define Q_λ to be the closure of the idempotent E_λ of R_n^n where n is the number of cells of λ ,

$$Q_\lambda = \Delta_n(E_\lambda) \in \mathcal{C}.$$

By looking at the limit $s \rightarrow 1$ it is straightforward to verify that the Homfly polynomial of Q_λ as an element of the plane is non-zero and therefore Q_λ is non-zero in $\mathcal{C}$ (see [7]).

We define a linear map Γ from $\mathcal{C}_+$ to $\mathcal{C}_+$ that is the encircling of a diagram in the annulus by a single loop with a specified orientation as shown in figure 4.

In R_n^n , one can write the identity braid on n strings with an encircling loop as the linear combination of so-called Jucys-Murphy operators as explained in [10]. The skein calculations in [2] then immediately imply the following lemma.

Lemma 4.1 *For any Young diagram λ we have $\Gamma(Q_\lambda) = c_\lambda Q_\lambda$ in $\mathcal{C}_+$ with the scalar*

$$c_\lambda = \frac{v^{-1} - v}{s - s^{-1}} + vs^{-1} \sum_{k=1}^{l(\lambda)} (s^{2(k-\lambda_k)} - s^{2k}).$$

Remark It is easy to confirm that the eigenvalues c_λ are pairwise different for all Young diagrams λ .

Lemma 4.2 *The set $\{Q_\lambda \mid \lambda \text{ has } n \text{ cells}\}$ is a basis for $\mathcal{C}_n$ for any $n \geq 0$.*

Proof We define $X_i^+ \in \mathcal{C}_i$ as the closure of the braid $\sigma_{i-1}\sigma_{i-2}\cdots\sigma_1 \in R_i^i$. The *weighted degree* of a monomial $(X_{i_1}^+)^{j_1}\cdots(X_{i_k}^+)^{j_k}$ is defined as $i_1j_1 + \cdots + i_kj_k$.

Any diagram in the annulus can be written inductively via the skein relations as a linear combination of totally descending curves. This means that $\mathcal{C}_n$ is spanned linearly by the monomials in the X_i^+ of weighted degree n . Hence, the dimension of $\mathcal{C}_n$ is at most $\pi(n)$, by which we denote the number of partitions of n .

On the other hand, Q_λ lies in $\mathcal{C}_n$ provided that the Young diagram λ has n cells. These elements are non-zero and they are linearly independent since they have pairwise different eigenvalues under the map Γ by lemma 4.1. By definition, there are $\pi(n)$ Young diagrams with n cells. Since the dimension of $\mathcal{C}_n$ is at most $\pi(n)$ by the above argument, the dimension of $\mathcal{C}_n$ is exactly $\pi(n)$ and the set of Q_λ where λ has n cells is a basis. $\diamond$

Remark Lemma 4.2 implies that the elements $X_i^+, i \geq 0$, generate $\mathcal{C}_+$ freely as a commutative algebra. This is a special case of Turaev's result [11] that $\mathcal{C}$ is generated freely as a commutative algebra by X_i^+, X_i^- for $i \geq 0$ where X_i^- derives from X_i^+ by reversing the orientation.

5 The variant skein $\mathcal{C}'$ of the annulus

The skein of the annulus $\mathcal{C}'$ with two boundary points has been considered e.g. in [6] and [4]. We are using the version introduced by Morton in [10] where the two boundary points lie on different boundary components of the annulus. This version has the benefit of a commutative multiplication.

We equip the annulus with two distinguished boundary points. An output point on the inner circle, and an input point on the outer circle. The resulting skein $\mathcal{C}'$ becomes an algebra by defining the product $A \cdot B$ of diagrams A and B as putting the outward circle of B next to the inward circle of A so that the two involved distinguished boundary points become a single point in the interior of the new annulus. An example for the multiplication is depicted in figure 5. The single straight arc e connecting the two marked points is the identity element, as shown in figure 6.

Any arc which connects the two marked points without crossings is regularly isotopic to a power of the diagram a which is depicted in figure 7.

For any integer $n \geq 1$ we have a linear map $\Delta'_n : R_n^n \rightarrow \mathcal{C}'$ as shown in figure 8. We abbreviate Δ'_n by Δ' if the index n is obvious from the context. With this notation, the diagram a from figure 7 is the image of the identity braid 1_2 of R_2^2 under the map Δ'_2 .

We denote the image of R_n^n under the map Δ'_n by $\mathcal{C}_n$. We define $\mathcal{C}'_+$ to be the submodule of $\mathcal{C}'$ spanned by all submodules $\mathcal{C}'_0, \mathcal{C}'_1, \dots$




Figure 5: The multiplication in $\mathcal{C}'$.




Figure 6: The identity e in $\mathcal{C}'$.




Figure 7: The arc a (at the left) and its inverse a^{-1} (at the right).




Figure 8: Map Δ'_n from R^n_n to $\mathcal{C}'$.




Figure 9: The additional arc for the closure.

The diagrams $A \cdot B$ and $B \cdot A$ are not regularly isotopic in general but they are equal modulo the skein relations as will be shown in lemma 5.1.

We note that $\mathcal{C}'$ can be turned into an algebra over $\mathcal{C}$ in two ways. Let A be a diagram in $\mathcal{C}'$ and γ be a diagram in $\mathcal{C}$. Then $\gamma \cdot A$ is defined as putting γ above A , and $A \cdot \gamma$ is defined as putting γ below A . Morton uses in [10] the notation $l(\gamma, A)$ resp. $r(A, \gamma)$ for the operation of $\mathcal{C}$ on $\mathcal{C}'$ on the left resp. right (up to a turn over of the annulus).

Lemma 5.1 ([10]) *$\mathcal{C}'$ is commutative.*

Proof Let D be a diagram in $\mathcal{C}'$. By induction on the number of crossings of D , we can write D as a scalar linear combination of diagrams, $D = \alpha_1 D_1 + \dots + \alpha_k D_k$, such that the single arc of any diagram D_i is totally descending and that it lies completely below any other component of D_i . Hence, $D = \gamma_1 a^{i_1} + \dots + \gamma_k a^{i_k}$ for elements $\gamma_1, \dots, \gamma_k$ of $\mathcal{C}$ and $i_j \in \mathbb{Z}$.

The commutativity of $\mathcal{C}'$ now follows from the observation that terms of the kind γa^i for $\gamma \in \mathcal{C}$ and $i \in \mathbb{Z}$ commute with each other. $\diamond$

Remark The above proof suggests that $\mathcal{C}'_+$ is the polynomial ring in a with coefficients in $\mathcal{C}_+$. This turns out to be true. One has to check that the powers of a are linearly independent over the scalars $\mathcal{C}_+$ (see [4] and [7]).

We define a closing operation $r \mapsto \hat{r}$ from $\mathcal{C}'$ to $\mathcal{C}$ which means adding the arc in figure 9 from above to a diagram r . In order that this is possible, the annulus for $\mathcal{C}$ has to be slightly larger than $\mathcal{C}'$. The framing of the diagram $\hat{r}$ is defined to be its blackboard framing. We remark that this closing operation is not an algebra homomorphism.

6 Basic skein relations in $\mathcal{C}'$

We recall that the quasi-idempotent $a_i \in R_i^i$ satisfies $a_i a_i = \alpha_i a_i$ for a non-zero scalar α_i . We define $h_i = \frac{1}{\alpha_i} \Delta(a_i) \in \mathcal{C}_i$ and $h'_i = \frac{1}{\alpha_i} \Delta'(a_i) \in \mathcal{C}'_i$ for any integer $i \geq 0$. We define $h_i = 0$ and $h'_i = 0$ for any integer $i < 0$.




Figure 10: Depiction of lemma 6.1.

The following lemma is a direct consequence of the decomposition

$$a_{i+1} = (a_i \otimes 1_1)(1_{i+1} + s\sigma_i + s^2\sigma_i\sigma_{i-1} + \cdots + s^i\sigma_i\sigma_{i-1} \cdots \sigma_1)$$

of the quasi-idempotent $a_{i+1} \in R_{i+1}^{i+1}$ (see [10], [7] for a proof). The lemma is depicted in figure 10.

Lemma 6.1 *We have*

$$[i+1]h'_{i+1} = eh_i + s^{-1}[i]h'_i a$$

in $\mathcal{C}'_+$ for any integer $i \geq 0$.

We define

$$t_i = h_i e - eh_i \in \mathcal{C}'_+$$

for any integer i . We remark that $t_i = 0$ for $i \leq 0$.

Lemma 6.2 *We have*

$$t_i = (s^{-i} - s^i)h'_i a$$

for any integer $i \geq 0$.

Proof We have

$$[i+1]h'_{i+1} = eh_i + s^{-1}[i]h'_i a$$

by lemma 6.1 for any integer i . By applying the map ρ from section 1 we get

$$[i+1]h'_{i+1} = h_i e + s[i]h'_i a$$

because h_i , h'_i and h'_{i+1} are invariant under ρ by lemma 3.2. These two equations imply that $t_i = h_i e - eh_i = (s^{-i} - s^i)h'_i a$. $\diamond$




Figure 11: The closure of $\Delta'(a_i)a$.

Corollary 6.3 *We have*

$$\hat{t}_i = (s^{1-2i} - s)vh_i$$

for any integer i .

Proof From lemma 6.2 and the skein relation in figure 11 we deduce that

$$\begin{aligned} \hat{t}_i &= (s^{-i} - s^i)vs^{1-i}h_i \\ &= (s^{1-2i} - s)vh_i \end{aligned}$$

for any integer $i \geq 0$. This equation holds for negative integers i as well because h_i and t_i are equal to zero for negative i . $\diamond$

Lemma 6.4 *We have*

$$\begin{vmatrix} t_i & t_{i+1} \\ t_j & t_{j+1} \end{vmatrix} = (s^2 - 1) \begin{vmatrix} eh_i & t_{i+1} \\ eh_j & t_{j+1} \end{vmatrix}$$

in $\mathcal{C}'$ for any integers i and j .

Proof If either i or j is negative then the lemma is obviously true because $t_k = 0$ for $k \leq 0$. Let $i \geq 0$ and $j \geq 0$ from now on. The combination of lemmas 6.1 and 6.2 implies that

$$[i+1]h'_{i+1} = eh_i + \frac{1}{1-s^2}t_i.$$

We have $(s^{-1} - s)[j+1]h'_{j+1}a = t_{j+1}$ by lemma 6.2. Multiplication of the above equation by these terms gives

$$(s^{-1} - s)[i+1][j+1]h'_{i+1}h'_{j+1}a = (eh_i)t_{j+1} + \frac{1}{1-s^2}t_it_{j+1}.$$

The left hand side of the above equation is invariant under the interchange of i and j because $\mathcal{C}'$ is commutative, and thus the right hand side is invariant under this interchange. Hence,

$$(eh_i)t_{j+1} + \frac{1}{1-s^2}t_it_{j+1} = (eh_j)t_{i+1} + \frac{1}{1-s^2}t_jt_{i+1}$$

which is equivalent to our claim. $\diamond$

Lemma 6.4 is the stepping stone from skein calculations to the following algebraic calculations.

7 Determinantal calculations in $\mathcal{C}'$

Lemma 7.1 *For any integer $r \geq 2$ and integers $i_1, i_2, \dots, i_r$ we have an equality of $(r \times r)$ -determinants in $\mathcal{C}'_+$*

$$\begin{vmatrix} h_{i_1}e & \cdots & h_{i_1+r-2}e & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ h_{i_r}e & \cdots & h_{i_r+r-2}e & t_{i_r+r-1} \end{vmatrix} = s^{2(r-1)} \begin{vmatrix} eh_{i_1} & \cdots & eh_{i_1+r-2} & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ eh_{i_r} & \cdots & eh_{i_r+r-2} & t_{i_r+r-1} \end{vmatrix}.$$

Proof Since $t_i = h_i e - e h_i$, we deduce from lemma 6.4 by the multilinearity of the determinant that

$$\begin{vmatrix} h_i e & t_{i+1} \\ h_j e & t_{j+1} \end{vmatrix} = s^2 \begin{vmatrix} e h_i & t_{i+1} \\ e h_j & t_{j+1} \end{vmatrix} \quad (7.1)$$

and

$$\begin{vmatrix} t_i & t_{i+1} \\ t_j & t_{j+1} \end{vmatrix} = (1 - s^{-2}) \begin{vmatrix} h_i e & t_{i+1} \\ h_j e & t_{j+1} \end{vmatrix}. \quad (7.2)$$

We remark that equation (7.1) is our claim in the case $r = 2$.

From now on let $r \geq 3$. We see that

$$\begin{vmatrix} t_{i_1} & t_{i_1+1} & t_{i_1+2} & \cdots & t_{i_1+r-1} \\ \vdots & \vdots & \vdots & & \vdots \\ t_{i_r} & t_{i_r+1} & t_{i_r+2} & \cdots & t_{i_r+r-1} \end{vmatrix} = (1 - s^{-2}) \begin{vmatrix} h_{i_1}e & t_{i_1+1} & t_{i_1+2} & \cdots & t_{i_1+r-1} \\ \vdots & \vdots & \vdots & & \vdots \\ h_{i_r}e & t_{i_r+1} & t_{i_r+2} & \cdots & t_{i_r+r-1} \end{vmatrix}$$

by developing the determinant on the left hand side by the first two columns, applying equation (7.2) to each summand, and redeveloping the determinant. By

doing this successively for the columns 1 and 2, 2 and 3, ..., $(r-1)$ and r , we deduce that

$$\begin{vmatrix} t_{i_1} & \cdots & t_{i_1+r-2} & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ t_{i_r} & \cdots & t_{i_r+r-2} & t_{i_r+r-1} \end{vmatrix} = (1-s^{-2})^{r-1} \begin{vmatrix} h_{i_1}e & \cdots & h_{i_1+r-2}e & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ h_{i_r}e & \cdots & h_{i_r+r-2}e & t_{i_r+r-1} \end{vmatrix}.$$

On the other hand, if we use the equation from lemma 6.4 instead of equation (7.2) in the above argument, we get

$$\begin{vmatrix} t_{i_1} & \cdots & t_{i_1+r-2} & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ t_{i_r} & \cdots & t_{i_r+r-2} & t_{i_r+r-1} \end{vmatrix} = (s^2-1)^{r-1} \begin{vmatrix} eh_{i_1} & \cdots & eh_{i_1+r-2} & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ eh_{i_r} & \cdots & eh_{i_r+r-2} & t_{i_r+r-1} \end{vmatrix}.$$

The above two equations imply that

$$\begin{vmatrix} h_{i_1}e & \cdots & h_{i_1+r-2}e & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ h_{i_r}e & \cdots & h_{i_r+r-2}e & t_{i_r+r-1} \end{vmatrix} = s^{2(r-1)} \begin{vmatrix} eh_{i_1} & \cdots & eh_{i_1+r-2} & t_{i_1+r-1} \\ \vdots & & \vdots & \vdots \\ eh_{i_r} & \cdots & eh_{i_r+r-2} & t_{i_r+r-1} \end{vmatrix}.$$

◇

The ring of symmetric functions (in countably many variables) is freely generated as a commutative algebra by the complete symmetric functions $h_1, h_2, \dots$. We therefore have a well defined map from the ring of symmetric functions to the skein of the annulus by mapping the complete symmetric function h_i to the skein element $h_i \in \mathcal{C}$. We denote the image of the Schur function s_λ by S_λ . The Schur function s_λ can be expressed in terms of the complete symmetric functions via the formula $s_\lambda = \det(h_{\lambda_i+j-i})_{1 \leq i, j \leq l(\lambda)}$ from [8]. We therefore have

$$S_\lambda = \det(h_{\lambda_i+j-i})_{1 \leq i, j \leq l(\lambda)} \in \mathcal{C}_n$$

where $n = |\lambda|$. We recall the linear map Γ from section 4.

Theorem 7.2 *We have $\Gamma(S_\lambda) = q_\lambda S_\lambda$ in $\mathcal{C}_+$ with the scalar*

$$q_\lambda = \frac{v^{-1} - v}{s - s^{-1}} + vs^{-1} \sum_{k=1}^{l(\lambda)} (s^{2(k-\lambda_k)} - s^{2k}).$$

Proof For any elements α and β of the skein of the annulus $\mathcal{C}$ we have $(\alpha e) \cdot (\beta e) = (\alpha \beta) e$ in $\mathcal{C}'$ where e is the identity of $\mathcal{C}'$. Hence

$$S_\lambda e = \det(h_{\lambda_i+j-i}e)_{1 \leq i, j \leq l(\lambda)}.$$

Similarly

$$eS_\lambda = \det(eh_{\lambda_i+j-i})_{1 \leq i,j \leq l(\lambda)}.$$

We denote $l(\lambda)$ by n from now on. We remark that the closure $(eS_\lambda)^\wedge$ is equal to S_λ with a disjoint circle which can be removed at the expense of the scalar $(v^{-1} - v)/(s - s^{-1})$. The closure $(S_\lambda e)^\wedge$ is equal to $\Gamma(S_\lambda)$.

By the multilinearity of the determinant we can write the difference of any two $(n \times n)$ -determinants as a telescope sum of n $(n \times n)$ -determinants.

$$\begin{aligned} & \begin{vmatrix} y_{11} & \cdots & y_{1n} \\ \vdots & & \vdots \\ y_{n1} & \cdots & y_{nn} \end{vmatrix} - \begin{vmatrix} z_{11} & \cdots & z_{1n} \\ \vdots & & \vdots \\ z_{n1} & \cdots & z_{nn} \end{vmatrix} = \\ & \sum_{k=1}^n \begin{vmatrix} y_{11} & \cdots & y_{1k-1} & (y_{1k} - z_{1k}) & z_{1k+1} & \cdots & z_{1n} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ y_{n1} & \cdots & y_{nk-1} & (y_{nk} - z_{nk}) & z_{nk+1} & \cdots & z_{nn} \end{vmatrix}. \end{aligned}$$

Applying this formula to the determinants for $S_\lambda e$ and eS_λ we get

$$S_\lambda e - eS_\lambda = \sum_{k=1}^n \begin{vmatrix} h_{\lambda_1} e & \cdots & h_{\lambda_1+k-2} e & t_{\lambda_1+k-1} & eh_{\lambda_1+k} & \cdots & eh_{\lambda_1+n-1} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ h_{\lambda_n+1-n} e & \cdots & h_{\lambda_n+k-1-n} e & t_{\lambda_n+k-n} & eh_{\lambda_n+k+1-n} & \cdots & eh_{\lambda_n} \end{vmatrix}.$$

By lemma 7.1 we deduce that

$$S_\lambda e - eS_\lambda = \sum_{k=1}^n s^{2(k-1)} \begin{vmatrix} eh_{\lambda_1} & \cdots & eh_{\lambda_1+k-2} & t_{\lambda_1+k-1} & eh_{\lambda_1+k} & \cdots & eh_{\lambda_1+n-1} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ eh_{\lambda_n+1-n} & \cdots & eh_{\lambda_n+k-1-n} & t_{\lambda_n+k-n} & eh_{\lambda_n+k+1-n} & \cdots & eh_{\lambda_n} \end{vmatrix}.$$

The n determinants appearing here are very special because each of them is a sum of terms of the form of a t_i above a product of h_j 's. Therefore the closure of each determinant is $\hat{t}_i$ above a product of h_j 's. Explicitly,

$$(S_\lambda e)^\wedge - (eS_\lambda)^\wedge = \sum_{k=1}^n s^{2(k-1)} \begin{vmatrix} h_{\lambda_1} & \cdots & h_{\lambda_1+k-2} & \hat{t}_{\lambda_1+k-1} & h_{\lambda_1+k} & \cdots & h_{\lambda_1+n-1} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ h_{\lambda_n+1-n} & \cdots & h_{\lambda_n+k-1-n} & \hat{t}_{\lambda_n+k-n} & h_{\lambda_n+k+1-n} & \cdots & h_{\lambda_n} \end{vmatrix}.$$

We know by corollary 6.3 that $\hat{t}_i$ is a scalar multiple of h_i . Hence

$$(S_\lambda e)^\wedge - (eS_\lambda)^\wedge = \sum_{k=1}^n \begin{vmatrix} h_{\lambda_1} & \cdots & h_{\lambda_1+k-2} & \beta_{1k} h_{\lambda_1+k-1} & h_{\lambda_1+k} & \cdots & h_{\lambda_1+n-1} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ h_{\lambda_n+1-n} & \cdots & h_{\lambda_n+k-1-n} & \beta_{nk} h_{\lambda_n+k-n} & h_{\lambda_n+k+1-n} & \cdots & h_{\lambda_n} \end{vmatrix}$$

where $\beta_{ik} = s^{2(k-1)}(s^{1-2(\lambda_i+k-i)} - s)v$. We use the notation $\rho_i = s^{2i-2\lambda_i-1}v$ and $\gamma_k = -s^{2k-1}v$, hence $\beta_{ik} = \rho_i + \gamma_k$. By the multilinearity of the determinant we get

$$(S_\lambda e)^\wedge - (eS_\lambda)^\wedge = (\gamma_1 + \cdots + \gamma_n)S_\lambda + \sum_{k=1}^n \begin{vmatrix} h_{\lambda_1} & \cdots & h_{\lambda_1+k-2} & \rho_1 h_{\lambda_1+k-1} & h_{\lambda_1+k} & \cdots & h_{\lambda_1+n-1} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ h_{\lambda_n+1-n} & \cdots & h_{\lambda_n+k-1-n} & \rho_n h_{\lambda_n+k-n} & h_{\lambda_n+k+1-n} & \cdots & h_{\lambda_n} \end{vmatrix}.$$

We bring the sum over the determinants in a more appropriate form via the general formula for variables w_{ij} and ρ_k ,

$$\sum_{k=1}^n \begin{vmatrix} w_{11} & \cdots & w_{1k-1} & \rho_1 w_{1k} & w_{1k+1} & \cdots & w_{1n} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ w_{n1} & \cdots & w_{nk-1} & \rho_n w_{nk} & w_{nk+1} & \cdots & w_{nn} \end{vmatrix} = (\rho_1 + \cdots + \rho_n) \begin{vmatrix} w_{11} & \cdots & w_{1n} \\ \vdots & & \vdots \\ w_{n1} & \cdots & w_{nn} \end{vmatrix}.$$

Applying this formula we get

$$\begin{aligned} (S_\lambda e)^\wedge - (eS_\lambda)^\wedge &= (\gamma_1 + \cdots + \gamma_n)S_\lambda + (\rho_1 + \cdots + \rho_n)S_\lambda \\ &= (\beta_{11} + \cdots + \beta_{nn})S_\lambda. \end{aligned}$$

Since $(eS_\lambda)^\wedge = (v^{-1} - v)/(s - s^{-1})S_\lambda$, as mentioned above, we deduce that $(S_\lambda e)^\wedge = q_\lambda S_\lambda$ with

$$\begin{aligned} q_\lambda &= \frac{v^{-1} - v}{s - s^{-1}} + \beta_{11} + \cdots + \beta_{nn} \\ &= \frac{v^{-1} - v}{s - s^{-1}} + vs^{-1} \sum_{k=1}^n (s^{2(k-\lambda_k)} - s^{2k}). \end{aligned}$$

◇

8 Proof that $S_\lambda = Q_\lambda$

We have to introduce some notation. Let λ be a Young diagram. A *standard tableau* of λ is a labelling of the cells of λ with the integers $1, 2, \dots, |\lambda|$ which is increasing in each row from left to right and in each column from top to bottom. The Young diagram underlying a standard tableau t is denoted by $\lambda(t)$. The number of different standard tableaux for a Young diagram λ is denoted by $d(\lambda)$.

Theorem 8.1 S_λ is equal to Q_λ for any Young diagram λ .

Proof Q_λ is non-zero in $\mathcal{C}$. Since the scalars q_λ and c_λ from theorem 7.2 and from lemma 4.1 are equal, we have that S_λ and Q_λ are eigenvectors with the same eigenvalue under the map Γ . Possibly, $S_\lambda = 0$. The set of Q_λ for all Young diagrams λ with n cells is a linear basis for $\mathcal{C}_n$ by lemma 4.2. Furthermore, the eigenvalues c_λ are pairwise different.

Hence, we deduce that S_λ is a scalar multiple of Q_λ for any Young diagram λ with n cells. This scalar is a rational function in x , v and s , and it is possibly equal to zero.

We denote the Young diagram consisting of a single cell by $\square$. We have that $S_\square = Q_\square$ is the single core circle of the annulus. Hence, $S_\square^n = Q_\square^n$ are n parallel copies of the core circle of the annulus.

Macdonald states in [8] example I.3.11 the multiplication rule for the Schur function s_λ with the first power sum p_1 (which is equal to $s_\square$). This is a special case of the Littlewood-Richardson rule for the multiplication of Schur functions. In this case, the rule is

$$s_\mu s_\square = \sum_{\substack{\mu \subset \eta \\ |\eta| = |\mu| + 1}} s_\eta.$$

By applying this successively we get

$$s_\square^n = \sum_{|\lambda|=n} d_\lambda s_\lambda$$

where d_λ is the number of standard tableaux of λ . Therefore,

$$S_\square^n = \sum_{|\lambda|=n} d_\lambda S_\lambda \tag{8.3}$$

in $\mathcal{C}_n$.

Blanchet describes in [3] an explicit semi-simple decomposition of R_n^n . The lemmas he uses for his construction are also contained in [2] as explained in

[7]. Generalizing results of Wenzl [12], Blanchet constructs elements $\alpha_t\beta_\tau \in R_n^n$ (which are denoted $\alpha_{t\tau}$ here) for standard tableaux t and τ with $\lambda(t) = \lambda(\tau)$. These elements satisfy $\alpha_{t\tau}\alpha_{s\sigma} = \delta_{\tau s}\alpha_{t\sigma}$ where $\delta_{\tau s}$ is the Kronecker-Delta, i.e. the elements $\alpha_{t\tau}$ multiply like matrix-units. An important observation is that the closure $\Delta(\alpha_{t\tau}) \in \mathcal{C}_+$ is equal to zero if $t \neq \tau$, and $\Delta(\alpha_{t\tau}) = Q_{\lambda(t)}$ if $t = \tau$. Furthermore, the identity braid $\text{id}_n \in R_n^n$ is equal to the sum of all α_{tt} , i.e.

$$\text{id}_n = \sum_{|\lambda(t)|=n} \alpha_{tt}.$$

We apply the closure map $\Delta : R_n^n \rightarrow \mathcal{C}_n$ to the above equation and we get that

$$Q_\square^n = \sum_{|\lambda|=n} d_\lambda Q_\lambda. \quad (8.4)$$

Since $S_\square = Q_\square$, equations (8.3) and (8.4) imply that

$$\sum_{|\lambda|=n} d_\lambda (Q_\lambda - S_\lambda) = 0.$$

Since any S_λ differs from Q_λ by a scalar, and $\{Q_\lambda \mid \lambda \text{ has } n \text{ cells}\}$ is a basis of $\mathcal{C}_n$ by lemma 4.2, we get that $Q_\lambda = S_\lambda$. $\diamond$

Theorem 8.1 together with the definition of S_λ implies that

Theorem 8.2 *The linear map induced by $s_\lambda \mapsto Q_\lambda$ from the ring of symmetric functions to the skein of the annulus $\mathcal{C}$ is a ring homomorphism.*

Acknowledgements The results are part of my PhD thesis. I thank my supervisor Prof. Hugh Morton for his guidance and help. The thesis was funded by the DAAD by a HSP-III-grant.

References

- [1] A. K. Aiston. *Skein theoretic idempotents of Hecke algebras and quantum group invariants*. PhD thesis, University of Liverpool, 1996.
- [2] A. K. Aiston and H. R. Morton. Idempotents of Hecke algebras of type A . *J. Knot Theory Ramif.*, 7(4):463–487, 1998.
- [3] C. Blanchet. Hecke algebras, modular categories and 3-manifolds quantum invariants. *Topology*, 39(1):193–223, 2000.
- [4] P. M. Gilmer and J. Zhong. On the Homflypt skein module of $S^1 \times S^2$. *math.GT/0007125*, 2000.
- [5] A. Gyoja. A q -analogue of Young symmetrizer. *Osaka J. Math.*, 23(4):841–852, 1986.
- [6] K. Kawagoe. On the skeins in the annulus and applications to invariants of 3-manifolds. *J. Knot Theory Ramif.*, 7(2):187–203, 1998.
- [7] S. G. Lukac. *Homfly skeins and the Hopf link*. PhD thesis, University of Liverpool, 2001.
- [8] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford University Press, second edition, 1995.
- [9] H. R. Morton. Invariants of links and 3-manifolds from skein theory and from quantum groups. In M. Bozhüyük, editor, *Topics in knot theory*, pages 107–156. Kluwer, 1993. NATO ASI Series C 399.
- [10] H. R. Morton. Skein theory and the Murphy operators. To appear in the Proceedings of Knots 2000, Korea. *math.GT/0102098*, 2001.
- [11] V. G. Turaev. The Conway and Kauffman modules of a solid torus. *Issled. Topol.*, 6:79–89, 1988.
- [12] H. Wenzl. Hecke algebras of type A_n and subfactors. *Invent. Math.*, 92(2):349–383, 1988.